



Your Health. Your Safety. Our Commitment.

Individual Client Risk Assessment Toolkit for Health Care Settings



Individual Client Risk Assessment Toolkit for Health Care Settings

Copyright © 2017

Product Code: VPRASEEN0417

Public Services Health and Safety Association (PSHSA)

4950 Yonge Street, Suite 1800

Toronto, Ontario M2N 6K1

Canada

Telephone: 416-250-2131

Fax: 416-250-7484

Toll Free: 1-877-250-7444

Web site: www.pshsa.ca

Connect with us:

 [@PSHSAca](https://twitter.com/PSHSAca)

Please note that all information provided in this toolkit is general in nature and may not be appropriate for particular situations or circumstances. The toolkit is not intended to provide legal advice or replace the Occupational Health and Safety Act (OHSA), its regulations or other relevant legislation that may apply to your work setting. Under no circumstances shall Public Services Health & Safety Association (PSHSA) be responsible for any damage or other losses resulting from reliance upon the information given to you, and all such liabilities are specifically disclaimed to the full extent permitted by law.

All material copyright 2017 Public Services Health & Safety Association. You may make no claim to copyright in any materials incorporating or derived from these materials.

All other rights reserved.

Terms of Use

By accessing or using these Public Services Health & Safety Association (PSHSA) resource materials, you agree to be bound by these terms and conditions.

Content: Although PSHSA endeavors to ensure that the information provided within these resource materials is as accurate, complete and current as possible, PSHSA makes no representations or warranties about the information, including in respect of its accuracy, completeness or currency. PSHSA assumes no responsibility for any loss or damage to you or any other person, howsoever caused, that is in any way related to the information found within these resource materials or your use of it.

Intent: The content within these resource materials is provided for educational and general informational purposes. It should not be considered as solicitation, endorsement, suggestion, advice or recommendation to use, rely on, exploit or otherwise apply such information or services.

Copyright: These resource materials and their content are protected by Canadian and international intellectual property laws, regulations, treaties and conventions. The content of this document, in whole or in part, may be reproduced without permission for non-commercial use only and provided that appropriate credit is given to PSHSA. No changes & / or modifications other than those required to reflect the utilizing organizations structure and terminology can be made to this document without written permissions from PSHSA. These Terms of Use must be retained and communicated in full on any permitted reproductions, disseminations and work products.

Other intellectual property rights: No permission is granted for the use of any other intellectual property right, including official marks or symbols, trademarks, logos, domain names or images.

Document Name: Individual Client Risk Assessment Toolkit V1.1 VPRASEEN0417

Product Code: VPRASEEN0417

Version Date: 2017.04.25

Introduction

About PSHSA

Public Services Health & Safety Association (PSHSA) provides occupational health and safety training and consulting services to various Ontario public sectors. These include healthcare, education, municipalities, public safety and First Nations communities.

As a funded partner of the Ministry of Labour (MOL), we work to prevent and reduce workplace injuries and occupational diseases by helping organizations adopt best practices and meet legislative requirements. To create safer workplaces, employers and employees must work together to identify potential hazards and eliminate or control risks before injuries and illnesses occur.

Workplace Violence in Healthcare

Violence in the workplace is a complex issue. It's also one of the top health and safety concerns facing Ontario's healthcare sector today. Research shows that workplace violence is three times more likely to occur among healthcare workers than any other occupation, including police officers and prison guards (International Council of Nurses, 2001; Kingma, 2001).

Each year, Ontario's Workplace Safety & Insurance Board (WSIB) allows more than 600 violence-related claims involving healthcare workers. While this number is alarming, many more cases go unreported (Findorff, Wall, & Gerberick, 2005). Healthcare staff work hard to keep others healthy and safe, yet their work can put them at risk and leave them with debilitating physical and psychological trauma.

Legislative changes in Ontario have broadened our awareness of workplace violence, and have strengthened our understanding that it cannot be considered part of the job. Under the law, employees have the right to a workplace that is safe and free of violence. Employers must ensure that risks are identified and that every reasonable precautions in the circumstances is taken to protect workers from harm.

The Five PSHSA Toolkits

PSHSA has created five toolkits to help healthcare organizations protect staff from workplace violence, and meet legal responsibilities for ensuring healthy and safe workplaces. The toolkits are:

1. Workplace Violence Risk Assessment (WPVRA)
2. Individual Client Risk Assessment (ICRA)
3. Flagging
4. Security
5. Personal Safety Response System (PSRS)

Acknowledgements

PSHSA acknowledges and appreciates the time and expertise of the many healthcare workers, organizations, frontline staff and labour unions that participated in the guidance and development of this toolkit.

Working Group Members

Name	Organization
Adam Nagler	Ministry of Health & Long-Term Care
Chris Day	County of Renfrew Paramedic Service
Danielle Baker	North Bay Regional Health Centre
Danielle Latulippe-Larmand	Ontario Nurses' Association
Heidi Birks	Registered Practical Nurses' Association of Ontario
Isabell Taylor	Richmond Terrace Nursing Home, Behavioural Supports Ontario
Jayne Brooks Keller	Chartwell Royal Oak Long-term Care Residence
Karen Heffernan	Ableliving Services – Thrive Group
Lisa McCaskell	Ontario Public Services Employees Union
Patti Boucher	Advanced Gerontological Education (AGE) Juravinski Research Centre
Waheed Butt	Peel Police

Toolkit development was led by Shelly Hurry and supported by Era Mae Ferron, both of PSHSA

Table of Contents


Terms of Use	i
Introduction.....	iii
About PSHSA	iii
Workplace Violence in Healthcare.....	iii
The Five PSHSA Toolkits	iv
Acknowledgements.....	iv
About the ICRA Toolkit	v
Summary of the Violence Assessment Tool (VAT)	3
Acute Care Violence Assessment Tool (VAT)	5
Long Term Care Violence Assessment Tool (VAT)	7
Community Care Violence Assessment Tool (VAT)	9
EMS Violence Assessment Tool (VAT).....	11
Appendix A: Sample Interventions	13
Physical Environment of Work	13
Work Practices.....	14
Staffing.....	15
Training.....	15
Appendix B: Sample Policy.....	17
Appendix C: VAT Pocket Card	27
Appendix D: VAT Poster	29
References	31

About the ICRA Toolkit

This toolkit addresses Individual Client Risk Assessment (ICRA).

The ICRA toolkit has been developed to help identify behaviours and triggers associated with increased risk of violence so prevention measures for staff and the client may be taken. The toolkit contains:

1. A Violence Assessment Tool to be completed by healthcare providers or managers / supervisors which includes risk indicators, a risk rating scale and a list of potential triggers
2. A list of suggested control interventions
3. A sample policy. The Sample Policy Includes
 - Purpose — what the ICRA is
 - Definitions — a glossary of key terms
 - Roles and responsibilities — who implements it
 - Procedures — how it's implemented
 - Communication / training — employee awareness, training, education and applied learning
 - Evaluation and continual improvement



All of these components can be downloaded individually from our website at <http://www.pshsa.ca/workplace-violence>



All clients, at first contact, must undergo an Individual Client Risk Assessment.

Please note that this assessment:

- must be part of a complete risk assessment system that includes policy / procedures, training, communication and evaluation — please see the [Sample Policy in Appendix B](#) for details.
- may be easily integrated into new or existing violence prevention protocols.
- should be used in conjunction with a flagging and care planning system when a history of violent behaviour or increased risk of violence is identified.

Summary of the Violence Assessment Tool (VAT)

What

The Violence Assessment Tool provides a snapshot of a client's immediate risk of violence. With this insight, your healthcare team can efficiently assess the risk, apply control interventions if needed, and improve worker safety while helping to increase quality of care. The VAT can also help identify and streamline referrals to internal / external Behavioural Supports Ontario (BSO) staff for further assessment, care planning solutions and resources for older adults with cognitive impairments presenting with at-risk behaviours.

Following extensive stakeholder consultation, the VAT was adapted from the Broset Violence Checklist and the Dynamic Appraisal of Situational Aggression instrument for use in multiple care settings such as Mental Health/Addiction.

The VAT contains three sections:

1. Risk Indicators
 - History of Violence
2. Behaviours observed
3. Overall risk rating
4. Triggers / Contributing factors

Why

Under the law, employees have the right to be told about risks of harm and how to work safely. Employers and supervisors must ensure that risks are identified and “take every precaution reasonable in the circumstances for the protection of a worker” [25(2)(h); 27(2)(c)]. Taking precautions to control identified risks and keeping workers informed is central to the prevention and management of workplace violence in Ontario workplaces [32.0.3; 32.0.5]. This tool was designed to be used in conjunction with [PSHSA's Flagging Resources](#) and to meet legislative requirements under OSHA.

Where

The VAT is for use in acute care, long-term care, community care, and emergency services.

When

The VAT should be completed at first contact with the client, and according to your organization's policies and procedures (e.g., at triage in acute care and once every shift thereafter; in long-term care it might be used during pre-screening, admission, *BSO meeting* and between prescribed Ministry documentation such as *RAI-MDS -Resident Assessment Instrument- Minimum Data Set; at the start of each community care visit*). Depending on the client's individual circumstances,

Individual Client Risk Assessment (ICRA) Toolkit

further assessment may be required, particularly when history of violence is known or violent, aggressive, or responsive behaviour is observed.


How

In Section A, read the list of behaviours. Score 1 for a history of violence and 1 for each behaviour observed. Add the scores — the maximum is 12. Next, consult the Risk Rating Scale in Section B to determine whether the client's risk level is low, moderate, high or very high. Each level provides cues for further action to consider. If moderate, high or very high risk is determined, complete Section C to identify factors that may trigger or escalate violent, aggressive, or responsive behaviour. It can be helpful to involve the client and / or substitute decision maker identify triggers and prevention / safety measures and procedures.

Use the information collected and the intervention resources listed below to develop a violence prevention care plan and safety measures that will also protect workers.

Intervention Resources:

- [Appendix A – Sample interventions](#) (See Page 13)
- [Appendix B – Sample organizational policy](#) (See Page 17)



For Additional PSHSA Resources, visit the PSHSA website at website at <http://www.pshsa.ca/workplace-violence/>

- Flagging handbook
- Security toolkit
- Personal Safety Response System (PSRS) toolkit
- Assessing Violence in the Community: A Handbook for the Workplace

Acute Care Violence Assessment Tool (VAT)

This form is to be completed by clinical healthcare worker or manager/supervisor.

Patient’s Name: _____

Identification #: _____

☐ Initial Assessment

☐ Reassessment

Section A: Risk Indicators

Read the list of behaviours below and identify behaviours that will require specific care interventions. A score of 1 is applied for past occurrence of any of the History of Violence behaviours; and additional scores of 1 are applied for each observed behavior. Add the scores — the maximum is 12.

HISTORY OF VIOLENCE:	SCORE
Score 1 for past occurrence of any of the following:	
<div><div></div><div>Exercising physical force, in any setting, towards any person including a caregiver that caused or could have caused injury</div><div></div><div>Attempting to exercise physical force, in any setting, towards any person including a caregiver that could cause injury</div><div></div><div>Statement or behaviours that could reasonably be interpreted as threatening to exercise physical force, in any setting, against any person including a caregiver that could cause injury</div></div>	
OBSERVED BEHAVIORS:	SCORE
Score 1 for each of the observed behaviour categories below.	
Confused (Disoriented – e.g., unaware of time, place, or person)	
Irritable (Easily annoyed or angered; Unable to tolerate the presence of others; Unwilling to follow instructions)	
Boisterous (Overly loud or noisy – e.g., slamming doors, shouting etc.)	
Verbal Threats (Raises voice in an intimidating or threatening way; Shouts angrily, insulting others or swearing; Makes aggressive sounds)	
Physical Threats (Raises arms / legs in an aggressive or agitated way; Makes a fist; Takes an aggressive stance; Moves / lunges forcefully towards others)	
Attacking Objects (Throws objects; Bangs or breaks windows; Kicks object; Smashes furniture)	
Agitate/Impulsive (Unable to remain composed; Quick to overreact to real and imagined disappointments; Troubled, nervous, restless or upset; Spontaneous, hasty, or emotional)	
Paranoid / suspicious (Unreasonably or obsessively anxious; Overly suspicious or mistrustful – e.g., belief of being spied on or someone conspiring to hurt them)	
Substance intoxication / withdrawal (Intoxicated or in withdrawal from alcohol or drugs)	
Socially inappropriate / disruptive behaviour (Makes disruptive noises; Screams; Engages in self-abusive acts, sexual behaviour or inappropriate behaviour – e.g., hoarding, smearing feces / food, etc.)	
Body Language (Torso shield – arms / objects acting as a barrier; Puffed up chest – territorial dominance; Deep breathing / panting; Arm dominance – arms spread, behind head, on hips; Eyes – pupil dilation / constriction, rapid blinking, gazing; Lips – compression, sneering, blushing / blanching)	
TOTAL SCORE	
Patient’s Risk Rating: <input type="checkbox"/> Low (0) <input type="checkbox"/> Moderate (1-3) <input type="checkbox"/> High (4-5) <input type="checkbox"/> Very High (6+)	

Completed By (Name/ Designation) _____ Date: _____

Section B: Overall Risk Rating

Apply the total behaviour score to the Risk Rating Scale to determine whether the patient’s risk level is low, moderate, high or very high. Each level provides cues for further action to consider. If moderate or high / very high risk is determined, complete Section C to identify factors that may trigger or escalate violent, aggressive, or responsive behaviour and ensure the care plan includes measures to avoid or reduce risk behaviours identified.

Overall Score	Actions to take
Low Score of 0	<ul style="list-style-type: none">Continue to monitor and remain alert for any potential increase in riskCommunicate any change in behaviours, that may put others at risk to the unit manager / supervisorEnsure communication devices / processes are in place (e.g., phone, personal safety alarm, check-in protocol and / or global positioning tracking system)
Moderate Score of 1-3	<ul style="list-style-type: none">Apply flag alertPromptly notify manager / supervisor so they can inform relevant staff and coordinate appropriate patient placement, unit staffing, and workflowAlert security and request assistance as needed. Ensure to inform security of risk management planScan environment for potential risks and remove if possibleEnsure section c is completed and initiate the violence prevention care planning process– care plan should address known triggers, behaviours and include safety measures appropriate for the situation for both patients and workersUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered care)Be prepared to apply behaviour management and self-protection teachings according to organizational policy/ procedures that are appropriate for the situation - training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe MORB training, self-defenseEnsure communication devices / processes are in place (e.g., phone, personal safety alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk to the unit manager / supervisorInform client of vat results, when safe to do so otherOther: _____
High Score of 4-5 OR Very High Score of 6+	<ul style="list-style-type: none">Apply flag alertPromptly notify manager / supervisor so they can ensure relevant staff are on high alert and prepared to respondAlert security and request security assistance as needed. Ensure to inform security of risk management planScan environment for potential risks and remove if possibleEnsure section c is completed and initiate the violence prevention care planning process – care plan should address known triggers, behaviours and include safety measures appropriate for the situation for both patients and workersUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered care)Be prepared to apply behaviour management and self-protection teaching appropriate for the situation in accordance to organizational policy / procedures – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe, MORB training, self-defenseInitiate applicable referralsEnsure communication devices / processes are in place (e.g. Phone, personal safety alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to the unit manager / supervisor so they can coordinate appropriate patient placement, unit staffing, and workflowCall 911 / initiate code white response as necessaryInform client of vat results, when safe to do soOther: _____

Section C: Contributing Factors

Physical, psychological, environmental, and activity triggers can lead to or escalate violent, aggressive or responsive behaviours. Documenting known triggers and behaviours and asking your patient or substitute decision maker (SDM) to help identify them can help you manage them more effectively and safely. Use the information collected and the intervention resources listed on p.2 and p.11 to develop an individualized violence prevention care plan and a safety plan to protect workers at risk.

QUESTION FOR CLIENT:		CONSIDERATIONS – Select any that Apply		
To help us provide the best care possible, please describe if there is anything during your stay that could cause you to become agitated, upset or angry e.g., I am agitated when...	PHYSICAL <input type="checkbox"/> hunger <input type="checkbox"/> pain <input type="checkbox"/> infection <input type="checkbox"/> new medication <input type="checkbox"/> other_____	PSYCHOLOGICAL <input type="checkbox"/> fear <input type="checkbox"/> uncertainty <input type="checkbox"/> feeling neglected <input type="checkbox"/> loss of control <input type="checkbox"/> being told to calm down <input type="checkbox"/> being lectured <input type="checkbox"/> other_____	ENVIRONMENTAL <input type="checkbox"/> noise <input type="checkbox"/> lighting <input type="checkbox"/> temperature <input type="checkbox"/> scents <input type="checkbox"/> privacy <input type="checkbox"/> time of day <input type="checkbox"/> days of the week <input type="checkbox"/> visitors <input type="checkbox"/> small spaces/ overcrowding <input type="checkbox"/> other_____	ACTIVITY <input type="checkbox"/> bathing <input type="checkbox"/> medication <input type="checkbox"/> past experiences <input type="checkbox"/> toileting <input type="checkbox"/> changes in routine <input type="checkbox"/> resistance to care <input type="checkbox"/> other_____
	What works to prevent or reduce the behaviour(s) e.g., When I am agitated, it helps if I...	POTENTIAL DE-ESCALATION TECHNIQUES Identify potential de-escalation strategies using above information such as respect personal space, actively listen, offer choices, give eye contact, use humor		
<input type="checkbox"/> Go for a walk <input type="checkbox"/> Listen to music <input type="checkbox"/> Watch TV <input type="checkbox"/> Draw <input type="checkbox"/> Read (Bible/Book) <input type="checkbox"/> Have space and time alone <input type="checkbox"/> Talk 1:1 with_____ (who?) <input type="checkbox"/> Participate in activities <input type="checkbox"/> Consult a family member or friend				

Long Term Care Violence Assessment Tool (VAT)

This form is to be completed by clinical healthcare worker or manager/supervisor.

Resident’s Name: _____

Identification #: _____

Click on the box above to insert your logo

☐ Initial Assessment

☐ Reassessment

Section A: Risk Indicators

Read the list of behaviours below and identify behaviours that will require specific care interventions. A score of 1 is applied for past occurrence of any of the History of Violence behaviours; and additional scores of 1 are applied for each observed behavior. Add the scores — the maximum is 12.

HISTORY OF VIOLENCE: Score 1 for past occurrence of any of the following:	SCORE
<ul style="list-style-type: none">▪ Exercising physical force, in any setting, towards any person including a caregiver that caused or could have caused injury▪ Attempting to exercise physical force, in any setting, towards any person including a caregiver that could cause injury▪ Statement or behaviours that could reasonably be interpreted as threatening to exercise physical force, in any setting, against any person including a caregiver that could cause injury	
OBSERVED BEHAVIORS: Score 1 for each of the observed behaviour categories below.	SCORE
Confused (Disoriented – e.g., unaware of time, place, or person)	
Irritable (Easily annoyed or angered; Unable to tolerate the presence of others; Unwilling to follow instructions)	
Boisterous (Overly loud or noisy – e.g., slamming doors, shouting etc.)	
Verbal Threats (Raises voice in an intimidating or threatening way; Shouts angrily, insulting others or swearing; Makes aggressive sounds)	
Physical Threats (Raises arms / legs in an aggressive or agitated way; Makes a fist; Takes an aggressive stance; Moves / lunges forcefully towards others)	
Attacking Objects (Throws objects; Bangs or breaks windows; Kicks object; Smashes furniture)	
Agitate/Impulsive (Unable to remain composed; Quick to overreact to real and imagined disappointments; Troubled, nervous, restless or upset; Spontaneous, hasty, or emotional)	
Paranoid / suspicious (Unreasonably or obsessively anxious; Overly suspicious or mistrustful – e.g., belief of being spied on or someone conspiring to hurt them)	
Substance intoxication / withdrawal (Intoxicated or in withdrawal from alcohol or drugs)	
Socially inappropriate / disruptive behaviour (Makes disruptive noises; Screams; Engages in self-abusive acts, sexual behaviour or inappropriate behaviour – e.g., hoarding, smearing feces / food, etc.)	
Body Language (Torso shield – arms / objects acting as a barrier; Puffed up chest – territorial dominance; Deep breathing / panting; Arm dominance – arms spread, behind head, on hips; Eyes – pupil dilation / constriction, rapid blinking, gazing; Lips – compression, sneering, blushing / blanching)	
TOTAL SCORE	
Resident’s Risk Rating: <input type="checkbox"/> Low (0) <input type="checkbox"/> Moderate (1-3) <input type="checkbox"/> High (4-5) <input type="checkbox"/> Very High (6+)	

Completed By (Name/ Designation) _____ Date: _____

Section B: Overall Risk Rating

Apply the total behaviour score to the Risk Rating Scale to determine whether the resident’s risk level is low, moderate, high or very high. Each level provides cues for further action to consider. If moderate or high / very high risk is determined, complete Section C to identify factors that may trigger or escalate violent, aggressive, or responsive behaviour and ensure the care plan includes measures to avoid or reduce risk behaviours identified.

Overall Score	Actions to take
Low Score of 0	<ul style="list-style-type: none">Continue to monitor and remain alert for any potential increase in riskCommunicate any change in behaviours, that may put others at risk, to the unit manager / supervisorEnsure communication devices / processes are in place (e.g. Phone, personal safety alarm, check-in protocol and / or global positioning tracking system)
Moderate Score of 1-3	<ul style="list-style-type: none">Apply flag alertPromptly notify shift supervisor so they can inform relevant staff and coordinate appropriate resident placement, unit staffing, and workflowAlert back-up staff / security / or police and request assistance, when neededScan environment for potential risks and remove if possibleEnsure section c is completed and initiate the violence prevention care planning process – care plan should address known triggers, behaviours and include safety measures appropriate for the situation for both residents and workersUse effective therapeutic communication (e.g. Maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered careBe prepared to be prepared to apply behaviour management and self-protection teachings according to organizational policy/ procedures that are appropriate for the situation – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe MORB training, self-defenseCollaborate with Behavioural Support Ontario (BSO) trained staff / psychogeriatric resource consultant as requiredEnsure communication devices / processes are in place (e.g., phone, personal safety alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to the shift supervisorInform client or SDM of VAT results, when safe to do soOther: _____
High Score of 4-5 OR Very High Score of 6+	<ul style="list-style-type: none">Apply flag alertPromptly notify shift supervisor so they can ensure relevant staff are on high alert and prepared to respondAlert back-up staff / security /police and request assistance when neededScan environment for potential risks and remove if possibleEnsure section c is completed and initiate the violence prevention care planning process– care plan should address known triggers, behaviours and include safety measures appropriate for the situation for both residents and workersUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered careBe prepared to apply behaviour management and self-protection teaching appropriate for the situation in accordance to organizational policy / Montessori – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe, MORB training, self-defenseInitiate applicable referralsCollaborate with Behavioural Support Ontario (BSO) trained staff / psychogeriatric resource consultant as requiredEnsure communication devices / processes are in place (e.g., phone, personal safety alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to the unit manager / supervisorCall 911 / initiate code white response as necessaryInform client of vat results, when safe to do soOther: _____


Section C: Contributing Factors

Physical, psychological, environmental, and activity triggers can lead to or escalate violent, aggressive or responsive behaviours. Documenting known triggers and behaviours and asking your resident or substitute decision maker (SDM) to help identify them can help you manage them more effectively and safely. Use the information collected and the intervention resources listed on p.2 and p.11 to develop an individualized violence prevention care plan and a safety plan to protect workers at risk.

QUESTION FOR RESIDENT:		CONSIDERATIONS – Select any that Apply		
To help us provide the best care possible, please describe if there is anything during your stay that could cause you to become agitated, upset or angry e.g., I am agitated when...	PHYSICAL <input type="checkbox"/> hunger <input type="checkbox"/> pain <input type="checkbox"/> infection <input type="checkbox"/> new medication <input type="checkbox"/> other_____	PSYCHOLOGICAL <input type="checkbox"/> fear <input type="checkbox"/> uncertainty <input type="checkbox"/> feeling neglected <input type="checkbox"/> loss of control <input type="checkbox"/> being told to calm down <input type="checkbox"/> being lectured <input type="checkbox"/> other_____	ENVIRONMENTAL <input type="checkbox"/> noise <input type="checkbox"/> lighting <input type="checkbox"/> temperature <input type="checkbox"/> scents <input type="checkbox"/> privacy <input type="checkbox"/> time of day <input type="checkbox"/> days of the week <input type="checkbox"/> visitors <input type="checkbox"/> small spaces/ overcrowding <input type="checkbox"/> other_____	ACTIVITY <input type="checkbox"/> bathing <input type="checkbox"/> medication <input type="checkbox"/> past experiences <input type="checkbox"/> toileting <input type="checkbox"/> changes in routine <input type="checkbox"/> resistance to care <input type="checkbox"/> other_____
	What works to prevent or reduce the behaviour(s) e.g., When I am agitated, it helps if I...	POTENTIAL DE-ESCALATION TECHNIQUES Identify potential de-escalation strategies using above information such as respect personal space, actively listen, offer choices, give eye contact, use humor		
<input type="checkbox"/> Go for a walk <input type="checkbox"/> Listen to music <input type="checkbox"/> Watch TV <input type="checkbox"/> Draw <input type="checkbox"/> Read (Bible/Book) <input type="checkbox"/> Have space and time alone <input type="checkbox"/> Talk 1:1 with _____ (who?) <input type="checkbox"/> Participate in activities <input type="checkbox"/> Consult a family member or friend				

Community Care Violence Assessment Tool (VAT)

This form is to be completed by clinical healthcare worker or manager/supervisor.



Click on the box above to insert your logo

Client’s Name: _____

Identification #: _____

☐ Initial Assessment

☐ Reassessment

Section A: Risk Indicators

Read the list of behaviours below and identify behaviours that will require specific care interventions. A score of 1 is applied for past occurrence of any of the History of Violence behaviours; and additional scores of 1 are applied for each observed behavior. Add the scores — the maximum is 12.

HISTORY OF VIOLENCE:	SCORE
Score 1 for past occurrence of any of the following:	
<ul style="list-style-type: none">▪ Exercising physical force, in any setting, towards any person including a caregiver that caused or could have caused injury▪ Attempting to exercise physical force, in any setting, towards any person including a caregiver that could cause injury▪ Statement or behaviours that could reasonably be interpreted as threatening to exercise physical force, in any setting, against any person including a caregiver that could cause injury	
OBSERVED BEHAVIORS:	SCORE
Score 1 for each of the observed behaviour categories below.	
Confused (Disoriented – e.g., unaware of time, place, or person)	
Irritable (Easily annoyed or angered; Unable to tolerate the presence of others; Unwilling to follow instructions)	
Boisterous (Overly loud or noisy – e.g., slamming doors, shouting etc.)	
Verbal Threats (Raises voice in an intimidating or threatening way; Shouts angrily, insulting others or swearing; Makes aggressive sounds)	
Physical Threats (Raises arms / legs in an aggressive or agitated way; Makes a fist; Takes an aggressive stance; Moves / lunges forcefully towards others)	
Attacking Objects (Throws objects; Bangs or breaks windows; Kicks object; Smashes furniture)	
Agitate/Impulsive (Unable to remain composed; Quick to overreact to real and imagined disappointments; Troubled, nervous, restless or upset; Spontaneous, hasty, or emotional)	
Paranoid / suspicious (Unreasonably or obsessively anxious; Overly suspicious or mistrustful – e.g., belief of being spied on or someone conspiring to hurt them)	
Substance intoxication / withdrawal (Intoxicated or in withdrawal from alcohol or drugs)	
Socially inappropriate / disruptive behaviour (Makes disruptive noises; Screams; Engages in self-abusive acts, sexual behaviour or inappropriate behaviour – e.g., hoarding, smearing feces / food, etc.)	
Body Language (Torso shield – arms / objects acting as a barrier; Puffed up chest – territorial dominance; Deep breathing / panting; Arm dominance – arms spread, behind head, on hips; Eyes – pupil dilation / constriction, rapid blinking, gazing; Lips – compression, sneering, blushing / blanching)	
TOTAL SCORE	
Client’s Risk Rating: <input type="checkbox"/> Low (0) <input type="checkbox"/> Moderate (1-3) <input type="checkbox"/> High (4-5) <input type="checkbox"/> Very High (6+)	

Completed By (Name/ Designation) _____ Date: _____

Section B: Overall Risk Rating

Apply the total behaviour score to the Risk Rating Scale to determine whether the client’s risk level is low, moderate, high or very high. Each level provides cues for further action to consider. If moderate or high / very high risk is determined, complete Section C to identify factors that may trigger or escalate violent, aggressive, or responsive behaviour and ensure the care plan includes measures to avoid or reduce risk behaviours identified.

Overall Score	Actions to take
Low Score of 0	<ul style="list-style-type: none">Continue to monitor and remain alert for any potential increase in riskCommunicate any change in behaviours, that may put others at risk, to the unit manager / supervisorEnsure communication device / processes are in place – (e.g., phone, personal safety / man-down alarm, check-in protocol; respectfully terminate client engagement / visit if concerns arise)
Moderate Score of 1-3	<ul style="list-style-type: none">Apply flag alertPromptly notify program manager / supervisor so they can inform relevant staff and coordinate appropriate staffing, workflowAlert back-up staff / security / police and request assistance when neededScan environment for potential risks and remove if possibleArrange to meet client in a public location as neededEnsure section c is completed and initiate the violence prevention care planning process – care plan should address known triggers, behaviours and include safety measures appropriate for the situation for clients and workersUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered care.Be prepared to apply behaviour management and self-protection teachings appropriate for the situation in accordance to organizational policy / Montessori – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe, MORB training, self-defenseEnsure communication device / processes are in place – (e.g., phone, personal safety / man-down alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to manager / supervisorInform client or SDM of VAT results, when safe to do soOther: _____
High Score of 4-5 OR Very High Score of 6+	<ul style="list-style-type: none">Apply flag alertPromptly notify program manager / supervisor so they can ensure relevant staff are on high alert and prepared to respondAlert back-up staff / security / police and request assistance when neededScan environment for potential risks and remove if possibleArrange to meet client in a public location as neededEnsure section c is completed and initiate the violence prevention care planning process – care plan should address known triggers, behaviours and include safety measures appropriate for the situation for both clients and workersInitiate applicable referralsUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic, and provide person-centered careBe prepared to apply behaviour management and self-protection teachings appropriate for the situation in accordance to organizational policy / Montessori – training programs provided may include GPA, Montessori, SMG P.I.E.C.E.S, U-First, Stay Safe MORB training, self-defenseEnsure communication device / process is in place – (e.g., phone, personal safety / man-down alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to the program manager / supervisorCall 911 / activate PSRS as necessaryInform client of VAT results, when safe to do soOther: _____


Section C: Contributing Factors

Physical, psychological, environmental, and activity triggers can lead to or escalate violent, aggressive or responsive behaviours. Documenting known triggers and behaviours and asking your client or substitute decision maker (SDM) to help identify them can help you manage them more effectively and safely. Use the information collected and the intervention resources listed on p.2 and p.11 of the PSHSA Individual Client Risk Tool to develop an individualized violence prevention care plan and a safety plan to protect workers at risk.

QUESTION FOR CLIENT:		CONSIDERATIONS – Select any that Apply		
To help us provide the best care possible, please describe if there is anything during your stay that could cause you to become agitated, upset or angry e.g., I am agitated when...	PHYSICAL <input type="checkbox"/> hunger <input type="checkbox"/> pain <input type="checkbox"/> infection <input type="checkbox"/> new medication <input type="checkbox"/> other_____	PSYCHOLOGICAL <input type="checkbox"/> fear <input type="checkbox"/> uncertainty <input type="checkbox"/> feeling neglected <input type="checkbox"/> loss of control <input type="checkbox"/> being told to calm down <input type="checkbox"/> being lectured <input type="checkbox"/> other_____	ENVIRONMENTAL <input type="checkbox"/> noise <input type="checkbox"/> lighting <input type="checkbox"/> temperature <input type="checkbox"/> scents <input type="checkbox"/> privacy <input type="checkbox"/> time of day <input type="checkbox"/> days of the week <input type="checkbox"/> visitors <input type="checkbox"/> small spaces/ overcrowding <input type="checkbox"/> other_____	ACTIVITY <input type="checkbox"/> bathing <input type="checkbox"/> medication <input type="checkbox"/> past experiences <input type="checkbox"/> toileting <input type="checkbox"/> changes in routine <input type="checkbox"/> resistance to care <input type="checkbox"/> other_____
	What works to prevent or reduce the behaviour(s) e.g., When I am agitated, it helps if I...	POTENTIAL DE-ESCALATION TECHNIQUES Identify potential de-escalation strategies using above information such as respect personal space, actively listen, offer choices, give eye contact, use humor		

EMS Violence Assessment Tool (VAT)

This form is to be completed by clinical healthcare worker or manager/supervisor.



Click on the box above to insert your logo

Patient’s Name: _____

Identification #: _____

☐ Initial Assessment

☐ Reassessment

Section A: Risk Indicators

Read the list of behaviours below and identify behaviours that will require specific care interventions. A score of 1 is applied for past occurrence of any of the History of Violence behaviours; and additional scores of 1 are applied for each observed behavior. Add the scores — the maximum is 12.

HISTORY OF VIOLENCE: Score 1 for past occurrence of any of the following:	SCORE
<ul style="list-style-type: none">▪ Exercising physical force, in any setting, towards any person including a caregiver that caused or could have caused injury▪ Attempting to exercise physical force, in any setting, towards any person including a caregiver that could cause injury▪ Statement or behaviours that could reasonably be interpreted as threatening to exercise physical force, in any setting, against any person including a caregiver that could cause injury	
OBSERVED BEHAVIORS: Score 1 for each of the observed behaviour categories below if you see any of the following examples.	SCORE
Confused (Disoriented – e.g., unaware of time, place, or person)	
Irritable (Easily annoyed or angered; Unable to tolerate the presence of others; Unwilling to follow instructions)	
Boisterous (Overly loud or noisy – e.g., slamming doors, shouting etc.)	
Verbal Threats (Raises voice in an intimidating or threatening way; Shouts angrily, insulting others or swearing; Makes aggressive sounds)	
Physical Threats (Raises arms / legs in an aggressive or agitated way; Makes a fist; Takes an aggressive stance; Moves / lunges forcefully towards others)	
Attacking Objects (Throws objects; Bangs or breaks windows; Kicks object; Smashes furniture)	
Agitate/Impulsive (Unable to remain composed; Quick to overreact to real and imagined disappointments; Troubled, nervous, restless or upset; Spontaneous, hasty, or emotional)	
Paranoid / suspicious (Unreasonably or obsessively anxious; Overly suspicious or mistrustful – e.g., belief of being spied on or someone conspiring to hurt them)	
Substance intoxication / withdrawal (Intoxicated or in withdrawal from alcohol or drugs)	
Socially inappropriate / disruptive behaviour (Makes disruptive noises; Screams; Engages in self-abusive acts, sexual behaviour or inappropriate behaviour – e.g., hoarding, smearing feces / food, etc.)	
Body Language (Torso shield – arms / objects acting as a barrier; Puffed up chest – territorial dominance; Deep breathing / panting; Arm dominance – arms spread, behind head, on hips; Eyes – pupil dilation / constriction, rapid blinking, gazing; Lips – compression, sneering, blushing / blanching)	
TOTAL SCORE	
Patient’s Risk Rating: <input type="checkbox"/> Low (0) <input type="checkbox"/> Moderate (1-3) <input type="checkbox"/> High (4-5) <input type="checkbox"/> Very High (6+)	

Completed By (Name/ Designation) _____ Date: _____

Section B: Overall Risk Rating

Apply the total behaviour score to the Risk Rating Scale to determine whether the patient’s risk level is low, moderate, high or very high. Each level provides cues for further action to consider. If moderate or high / very high risk is determined, complete Section C to identify factors that may trigger or escalate violent, aggressive, or responsive behaviour and ensure the care plan includes measures to avoid or reduce risk behaviours identified.

Overall Score	Actions to take
Low Score of 0	<ul style="list-style-type: none">Continue to monitor and remain alert for any potential increase in riskCommunicate any change in behaviours, that may put others at risk to the manager / supervisorEnsure communication devices / processes are in place (e.g., radio, personal safety / man-down alarm, check-in protocol and / or global positioning tracking system)
Moderate Score of 1-3	<ul style="list-style-type: none">Apply flag alertPromptly notify manager / supervisor so they can inform relevant staff and coordinate appropriate staffing and workflowRequest additional assistance / police back-up, when neededScan environment for potential risks and remove if possibleAddress known triggers, behaviours and implement organization / provincially accepted safety measures appropriate for the situationUse effective therapeutic communication (e.g., maintain a calm, reassuring demeanor, remain non-judgmental and empathetic)Be prepared to apply behaviour management and self-protection techniques appropriate for the situation in accordance to organizational policy / Montessori – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe MORB training, self-defenseEnsure communication devices / processes are in place (e.g., radio, personal safety / man-down alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk to manager / supervisorInform client of VAT results, when safe to do so otherOther: _____
High Score of 4-5 OR Very High Score of 6+	<ul style="list-style-type: none">Apply flag alertPromptly notify program manager / supervisor so they can ensure relevant staff are on high alert and prepared to respondRequest additional assistance / police back-up, when neededScan environment for potential risks and remove if possibleAddress known triggers, behaviours and implement organization / provincially approved safety measures appropriate for the situation for both clients and workersInitiate applicable referralsUse effective therapeutic communication (maintain a calm, reassuring demeanor, remain non-judgmental and empathetic)Be prepared to apply behaviour management and self-protection techniques in accordance to organizational policy / Montessori – training programs provided may include GPA, Montessori, SMG, P.I.E.C.E.S, U-First, Stay Safe MORB training, self-defenseEnsure communication device / process is in place – (e.g., phone, personal safety / man-down alarm, check-in protocol and / or global positioning tracking system)Communicate any change in behaviours, that may put others at risk, to the program manager / supervisorActivate emergency communication codes as necessaryInform client of VAT results, when safe to do soOther: _____

Section C: Contributing Factors

Physical, psychological, environmental, and activity triggers can lead to or escalate violent, aggressive or responsive behaviours. Documenting known triggers and behaviours and asking your patient or substitute decision maker (SDM) to help identify them can help you manage them more effectively and safely. Use the information collected and the intervention resources listed on p.2 and p.11 of the PSHSA Individual Client Risk Tool to develop an individualized violence prevention care plan and a safety plan to protect workers at risk.

QUESTION FOR CLIENT:		CONSIDERATIONS – Select any that Apply		
To help us provide the best care possible, please describe if there is anything during your stay that could cause you to become agitated, upset or angry e.g., I am agitated when...	PHYSICAL <input type="checkbox"/> hunger <input type="checkbox"/> pain <input type="checkbox"/> infection <input type="checkbox"/> new medication <input type="checkbox"/> other_____	PSYCHOLOGICAL <input type="checkbox"/> fear <input type="checkbox"/> uncertainty <input type="checkbox"/> feeling neglected <input type="checkbox"/> loss of control <input type="checkbox"/> being told to calm down <input type="checkbox"/> being lectured <input type="checkbox"/> other_____	ENVIRONMENTAL <input type="checkbox"/> noise <input type="checkbox"/> lighting <input type="checkbox"/> temperature <input type="checkbox"/> scents <input type="checkbox"/> privacy <input type="checkbox"/> time of day <input type="checkbox"/> days of the week <input type="checkbox"/> visitors <input type="checkbox"/> small spaces/ overcrowding <input type="checkbox"/> other_____	ACTIVITY <input type="checkbox"/> bathing <input type="checkbox"/> medication <input type="checkbox"/> past experiences <input type="checkbox"/> toileting <input type="checkbox"/> changes in routine <input type="checkbox"/> resistance to care <input type="checkbox"/> other_____
	What works to prevent or reduce the behaviour(s) e.g., When I am agitated, it helps if I...	POTENTIAL DE-ESCALATION TECHNIQUES Identify potential de-escalation strategies using above information such as respect personal space, actively listen, offer choices, give eye contact, use humor		
<input type="checkbox"/> Go for a walk <input type="checkbox"/> Listen to music <input type="checkbox"/> Watch TV <input type="checkbox"/> Draw <input type="checkbox"/> Read (Bible/Book) <input type="checkbox"/> Have space and time alone <input type="checkbox"/> Talk 1:1 with _____ (who?) <input type="checkbox"/> Participate in activities <input type="checkbox"/> Consult a family member or friend				



Appendix A: Sample Interventions

Visit: pshsa.ca/workplace-violence

PHYSICAL ENVIRONMENT, WORK PRACTICES, STAFFING AND TRAINING

Most workplace-violence best practices categorize intervention controls into four main areas:

- Physical environment of work
- Work practices
- Staffing
- Training

Within these four areas, experts recommend the following hierarchy of control measures:

- Eliminate the hazard.
- Engineer solutions.
- Reorganize and provide training.
- Provide personal protective equipment.

Physical Environment of Work

- Create an environment that limits triggers of violent behaviours.
- Create an environment that reduces risk — e.g., good lighting, removal of hazardous furniture, and removal / securing of instruments that could be used as weapons.
 - Weighted or secured furniture
 - Rounded edges on furniture
 - Pictures with no frames or glass and secured to the wall
 - Secured access to staff rooms, kitchen areas
 - Access to needles or other medical equipment
 - Hallways, exit routes free from clutter
- Create an environment that promotes client comfort — e.g.,

Individual Client Risk Assessment (ICRA) Toolkit

minimize overcrowding, ensure comfortable seating, and provide visual displays or soothing music, distraction activities.

- Ensure balance between static security measures (e.g., enclosed reception area) and physical design (e.g., aesthetics value, home-like environment) is appropriate for the circumstance.
- Have a seclusion room available when needed.
- Implement security measures such as security guards and protective equipment (e.g. convex mirrors, personal alarms etc.).
 - Security must be well-trained on verbal de-escalation and relationship management.
 - See PSHSA's Security Toolkit as needed
- Implement security measures in the community — e.g., mobile phones, personal alarms and safety features in vehicles such as automatic locks and alarms.

Work Practices

- Complete VAT to ensure client information is accurate and risks, behaviours, triggers, and safety measures for staff and clients are documented.
- Ensure initial and ongoing assessment of clients.
- Assess the risk of potentially violent situations at every visit and assess them considering current client population, acuity, staffing and work flow.
- Implement procedures for organizational violence risk assessments to ensure information

is up-to-date – See [PSHSA's Workplace Violence Risk Assessment Toolkit](#) as needed.

- Implement procedures on care planning and behaviour planning.
- Implement measures and procedures to protect workers (e.g., Kevlar gloves, spit shields etc.)
- Implement flagging procedures that include identification of triggers, behaviours and safety measures for patients and workers – See [PSHSA's Flagging Handbook](#) as needed
- Use personal safety response systems — e.g., personal alarms and mobile phones.
- Adopt a buddy system.
- Establish and use Emergency Codes.
- Establish code Words (e.g., requesting a “Yellow card” to signal distress)
- Practice workplace violence emergency procedures through mock drills and scenario-based training.
- Adopt community care service agreement contracts that address violence.
- Establish communication procedures that ensure traceability of workers — e.g., check-in in / check-out practices.
- Implement BETSI — Behavioural Education and Training Supports Inventory (an education / training decision-making tool and program inventory)
- Conduct Critical Incident Debriefings.
- Develop policies/procedures for staff support post incident.

Staffing

- Ensure adequate staffing levels, skill set and competencies.
- Provide adequate training for staff, ensuring skills and experience meet client needs and protect workers' health and safety.
- Ensure staff responding to Emergency Codes (e.g., Code White, Code Purple, Code Silver) receive adequate education and training, developed in consultation with the JHSC or HSR that is appropriate to the highest level of response required (e.g., relationship management training, active listening, collaborative problem solving, self-defense).
- Allow for job rotation where possible in order to help reduce time in stressful working situations.
- Ensure sufficient security personnel, and that skills meet work demands.
- Adopt a buddy system to avoid having providers work alone with high-risk clients.
- Adopt a culture of trust, teamwork and support when a staff needs to relieve one another from a challenging or high-risk situation.

Training

- Ensure that, where necessary, staff receive the following training:
 - Corporate Workplace Violence and Harassment Policy
 - Individual Client Risk Assessment policy and procedures
 - Flagging and risk communication protocol
 - Behavioural Management Techniques (e.g., GPA, SMG, P.I.E.C.E.S, U-First Me & U-First, Stay Safe MORB training)
 - Self-defense, and Sharp-edged Weapons
 - Non-violent Crisis Intervention
 - Verbal De-escalation and Relationship Management
 - Emergency measures, Codes and responses
 - Security protocols (e.g., Access control, working alone, security guard / personnel functions-)
 - Personal safety alarms and summoning assistance
 - Use of restraints (e.g. restraint application on the floor) / seclusion
 - Reporting and investigating hazards and incidents
 - Stress Debriefing
 - Psychological Health & Safety in the Workplace



Appendix B: Sample Policy

Visit: pshsa.ca/workplace-violence

Sample Policy

Manual: Health and safety	Subject: Workplace Violence Client Risk Assessment	Policy number: XX-XXX-XX
Effective date: XX-XX-XXXX	Date revised: XX-XX-XXXX	Date of next review: XX-XX-XXXX
Policy reviewers:		
Authorized by:	Signature:	

About the ICRA Policy

This policy covers the following:

- **Purpose** — what the ICRA is
- **Statement** — sample policy wording
- **Definitions** — a glossary of key terms
- **Roles and responsibilities** — who implements it
- **Procedures** — how it's implemented
- **Communication / training** — employee orientation
- **Evaluation and continual improvement**

Scope

The ICRA tool was developed for employers and employees in the healthcare and emergency services sectors. It's designed to help these care providers identify risk factors and levels associated with workplace violence. By conducting regular client-risk assessments, providers can apply control interventions that promote both employee and client safety, as well as ensure client-centred care.

The ICRA should be completed at first contact with a client (e.g., triage) , and on an ongoing basis depending on client population, acuity levels, staffing, work flow, individual client circumstances, and the employer's operational policies and organizational risk assessment findings.

Objectives

Specific goals of the ICRA are to:

- Implement a practical, immediate and easy-to-use assessment tool that identifies a client's past history, observed behaviours, triggers, and risk factors associated with violence
- Facilitate early recognition of violence and enable early application of control interventions
- Identify client's overall levels of risk
- Help healthcare organizations develop effective prevention measures
- Establish control interventions for different client populations to manage moderate and high or very high-risk clients

Definitions

These behaviours, if understood, can be managed and prevented. Behavioural and environmental strategies play a crucial role in effectively managing violence and responsive behaviours.

Client

For the purpose of this tool, a client means a patient, resident, person that is being supported, a consumer, a family member / loved one, a visitor, or a police subject or accused.

Clinical healthcare worker

A clinical staff member who provides preventive, curative, promotional or rehabilitative healthcare services to clients.

First point of contact

For the purpose of this tool, first point of contact refers to the initial interaction with the clinical healthcare worker assessing client's care needs.

Flag

A visual and / or electronic alert used to inform staff of a risk of violent, aggressive or responsive behaviours and signal additional individualized care needs and preventive measures for staff and client.

Flagging

A standardized method to communicate safety-related concerns to workers.

Individual Client Risk Assessment (ICRA)

Systematic process used by healthcare professionals for evaluating a client's likelihood of violent, aggressive, or responsive behaviour.

Responsive Behaviours

A protective means by which persons with dementia or other conditions may communicate an unmet need (e.g., pain, cold, hunger, constipation, boredom) or reaction to their environment (e.g., lighting, noise, invasion of space).

Staffing

Staffing refers to the selection of workers required at various times and in various settings to ensure prevention or appropriate intervention. In a comprehensive workplace violence prevention program (WVPP), the staffing model should ensure that the required skill sets and core competencies are found within the interdisciplinary team at the point of care and at a broader organizational level. Required skill sets and core competencies may be identified through organizational and individual risk assessment processes and will vary from organization to organization.

Tool

For the purpose of this toolkit, a tool is an instrument — e.g., survey, guidelines, or checklist — that helps users accomplish a specific task that supports a specific evidence-based recommendation or practice standard.

Transition of care (TOC) / Transfer of accountability (TOA)

An interactive process for transferring client specific information from one healthcare worker to another or from one team of care providers to the next, to ensure continuity of care, as well as staff and client safety. Examples include:

- Nurse to nurse at change of shift
- Nurse to nurse when care is temporarily assigned to another nurse on a short term basis
- Transfer from one client care area to another
- When transferring to a different client care unit within the organization
- When transferring to an outside organization

Trigger

A circumstance / situation that impacts or escalates client's behaviour. Triggers may be physical, psychological, environmental, or activity-related.

Violent behaviour

Acts of violence including, but not limited to: choking, hitting, shoving, pushing, biting, spiting, shouting, swearing, verbal threats, groping, pinching, kicking, throwing objects, shaking fists, stabbing and threatening assault.

Workplace

Any land premises, location or thing at, upon, in or near which a worker works.

Workplace violence

It is defined by the Occupational Health and Safety Act as:

- the exercise of physical force by a person against a worker, in a workplace, that causes or could cause physical injury to the worker;
- an attempt to exercise physical force against a worker, in a workplace, that could cause physical injury to the worker; or,
- a statement or behaviour that it is reasonable for a worker to interpret as a threat to exercise physical force against the worker, in a workplace, that could cause physical injury to the worker.

There are four types of workplace violence:

- Type I (external perpetrator): The violent person has no relationship to the worker or workplace
- Type II (client or customer): The violent person is a client at the workplace who becomes violent toward a worker or another client
- Type III (employment-related): The violent person has / had some type of job-related involvement with the workplace.
- Type IV (domestic violence): The violent person has a personal relationship with an employee or a client

Roles & Responsibilities

The board of directors of an organization must take reasonable care to ensure the corporation complies with:

- The Occupational Health & Safety Act (OHSA) and its regulations
- Orders and requirements of inspectors and directors of the Ministry of Labour (MOL)
- Orders of the MOL

Employers are obligated to:

- Take every precaution reasonable in the circumstances for the protection of a worker
- Ensure the measures and procedures for the ICRA program are carried out
- Evaluate the effectiveness and use of the ICRA, in consultation with the organization's joint health and safety committee (JHSC) or health and safety (H&S) representative
- Develop, establish and deliver training and education for all employees on the use of the ICRA, in consultation with the JHSC or H&S representative
- Comply with the organization's workplace violence prevention program
- Comply with the organization's internal and external incident reporting obligations as outlined in the Incident Reporting and Investigation Policy and reporting requirements under the OHSA
- Comply with the organization's return-to-work program, as required
- Appoint competent supervisors
- Provide the JHSC with copies of all accident illness/reports as per OHSA requirements
- Provide JHSC with copies of all written risk assessments as per legislation

Managers / Supervisors are obligated to:

- Take every precaution reasonable in the circumstances for the protection of a worker
- Ensure the unit has resources to manage workplace violence that supports the number and risk level of clients identified at-risk.
- Ensure employees are trained on ICRA procedures and that new employees are trained at orientation.
- Provide refresher training to all employees at least once a year (or more often if required)
- Enforce the use of the ICRA and monitor worker compliance
- Monitor and evaluate the effectiveness of the ICRA through regular workplace inspections / audits
- Comply with the organization's workplace violence prevention program
- Comply with the organization's internal and external incident reporting obligations as outlined in the Incident Reporting and Investigation Policy and reporting requirements under the OHSA
- Comply with the organization's return-to-work program, as required

Employees are obligated to:

- Participate in education and training programs on the use of the ICRA
- Understand and comply with the use of the ICRA program
- Comply with the organization's workplace violence prevention program
- Comply with the organization's internal and external incident reporting obligations as outlined in the Incident Reporting and Investigation Policy
- Report all hazards to their supervisor
- Comply with the organization's return-to-work program, as required

The joint health and safety committee (JHSC) or health and safety representative (H&S representative) is obligated to:

- Ensure the employer has consulted about the development, establishment and implementation of the ICRA program
- Make recommendations to the employer for developing, establishing and providing ICRA training
- Comply with the organization's workplace violence prevention program
- Comply with the organization's internal and external incident reporting obligations as outlined in the Incident Reporting and Investigation Policy
- Be provided with copies of all accident illness/reports as per OHS requirements
- Be provided with copies of all risk assessments as per legislation
- Comply with the organization's return-to-work program, as required
- Gather feedback from employees during workplace inspections about assessment tool use, effectiveness, and suggested improvements.

Procedures

This list is not exhaustive nor is the organization required to use all measures and procedures listed. The organization must decide which measures and procedures best suit their operations and risk management/prevention needs to best protect workers and clients.

Prevention approach

Administer the ICRA at the first point of contact —e.g., at triage. For community care, Pre-visit and Pre-travel assessments should ideally be completed within 24 hours before the initial home visit, followed by a behaviour assessment such as VAT completed at the beginning of each home visit thereafter – refer to PSHSA's [Assessing Violence in the Community: A Handbook for the Workplace](#), as needed.

Continue to administer the ICRA at pre-determined times after initial contact as outlined in the organization's procedures — e.g., once every shift, weekly, or at the beginning of each home visit.

Individual Client Risk Assessment (ICRA) Toolkit

Recommended timing for different healthcare subsectors is as follows:

- Acute care/mental health/addictions — at first point of contact with a clinical healthcare worker (e.g., triage) and repeated during each shift as outlined by the organization.
- Long-term care:
 - at first point of contact with a clinical healthcare worker (e.g., upon admission or pre-assessment if applicable)
 - when client exhibits behaviours
 - between prescribed Ministry-required documentation such as the RAI-MDS (*Resident Assessment Instrument- Minimum Data Set*) or RAI-HC assessments (*Resident Assessment Instrument- Home Care*)
- Community care — upon contract acceptance; prior to the first home visit; at the start of each home visit
- Paramedics — at first point of contact, and prior to discharge to a healthcare provider
- Police — at first contact; prior to discharge to a healthcare provider; during hourly cell observation; or as outlined in organizational procedures

Repeat the ICRA process when a change in client behaviour warrants a reassessment.

Apply the Risk Rating Scale as part of the VAT— e.g., on every shift, prior to a home visit or at the start of a home visit. Note: This Risk Rating Scale is designed for use with this assessment tool only. If an organization chooses to use another ICRA tool, use the risk rating scale designed for that particular tool.

Scoring the Risk Rating Scale

The ICRA uses a series of questions to identify the presence of risk-related behaviours. A score of 1 is given for a history of violence and for observation of specific predetermined behaviours. The numbers are added to reach a total behaviour score. The score is then applied to the Risk Rating Scale to determine whether the client presents a low, moderate or high/very risk. Each risk level provides cues for further action to consider.

Client's Risk Rating: ☐ Low (0) ☐ Moderate (1-3) ☐ High (4-5) ☐ Very High (6+)

Protection approach

- Implement preventive measures for all moderate or high / very high risk clients, according to organizational policies and procedures. Use the intervention resources listed on page 4 and suggested measures outlined in [Appendix A](#) on page 13 as a guide.
- Adopt security measures and personal safety response systems (e.g., personal alarms, mobile phones) according to organizational policies and procedures.
- Develop a care plan or violence behaviour plan to identify, address, and minimize triggers.
- Develop a safety plan for all workers at risk
- Establish a strategy to communicate risk of workplace of violence, triggers, behaviours and prevention / safety measures. See [PSHSA's Flagging Handbook](#) as needed.

Post-incident response

- Apply organizational post-incident responses to reduce the negative impact of violence.
- Communicate debriefing results with all affected workers of violent incidents to reduce their negative impact in the workplace and prevent further incidents.

Reporting and investigation

- Refer to the organization's workplace violence prevention program for reporting and investigation procedures. Conduct and involve client or substitute decision maker in a root cause analysis to determine why client was triggered and develop/update care plan and worker safety measures.

Emergency response procedures

- Refer to the organization's emergency response procedures — e.g., Code White, staff alert, etc.

Transition of care / Transfer of accountability

- ICRA risk levels and recommended interventions should be communicated at all transitions of care.

Re-training

- A refresher on the use of the ICRA Tool is required annually, or more often/as outlined in the organization's policy. Large organizations should offer monthly or quarterly sessions to ensure all employees are informed.

Client Aggression Prevention Program training

- Refer to BETSI — Behavioural Education and Training Supports Inventory tool — to determine where training is needed.
- Promote a respectful workplace and communicate clear behaviour expectations for management, workers, physicians, contractors, clients and the general public.
- Focus on / review the organization's behaviour management program — e.g., Safe Management Group, P.I.E.C.E.S, Gentle Persuasive Approach, Crisis Prevention Institute, U-First Me & U-First, Stay Safe MORB training etc.
- Focus on / review the organization's Emergency Response, flagging, and security policies and procedures. See [PSHSA's Flagging Handbook](#), [Security Toolkit](#), and [PSRS Toolkit](#), as needed.

Communication / Training

- All applicable employees shall receive training/education on the organization's ICRA Policy. New employees will receive this training at orientation. Ongoing refresher training will be provided on a regular basis as part of routine violence-prevention training and when new procedures are developed or revised. The JHSC or HSR must be consulted in the development of such training.

Training should include:

- An understanding of violent, aggressive, and responsive behaviours at work
- Terminology around workplace violence and client aggression / responsive behaviours
- When and how often the ICRA tool is to be implemented
- How to determine level of risk
- How to choose appropriate control measures, as outlined in the organization's policies and procedures
- When and how to apply flag alerts
- Safety measures to protect workers and clients
- Security functions and protocols

Evaluation and Continual Improvement

To effectively evaluate the ICRA program, healthcare organizations should:

- Evaluate the program annually in consultation with the JHSC or H&S representative
- Evaluate the effectiveness of ICRA communication and training, using both leading and lagging indicators.

Individual Client Risk Assessment (ICRA) Toolkit

- Share findings with the JHSC or H&S representative and the board of directors

The organization's continual improvement plan should be supported by:

- root cause analysis
- corrective action points planned to resolution
- assigned responsibilities for each point
- expected timelines for each point
- adjust program and training based on evaluation

Organizations must monitor the plan regularly for compliance, ensuring supervisors are trained to support consistent use of the tools, communicate program outcomes, and follow-up on implementation challenges.



Appendix C: VAT Pocket Card

Visit: pshsa.ca/workplace-violence

Pocket Card Instructions

1. Download the template from www.pshsa.ca/workplace-violence
2. Print the template on a perforated card stock that contains 10 standard business card stock 2"x3.5" cards and works with Avery 5371, 5911, 8371 and 8859.

Pocket Card

<p>Before providing care, assess behaviours for potential risk</p> <ul style="list-style-type: none"> <input type="checkbox"/> History of Violence <input type="checkbox"/> Confused <input type="checkbox"/> Irritable <input type="checkbox"/> Boisterous <input type="checkbox"/> Verbal Threats <input type="checkbox"/> Physical Threats <input type="checkbox"/> Agitate/Impulsive <input type="checkbox"/> Paranoid/Suspicious <input type="checkbox"/> Substance <input type="checkbox"/> Intoxication/withdrawal <input type="checkbox"/> Socially appropriate/disruptive <input type="checkbox"/> Body Language <p>Each Check = 1 pt 0=Low Risk; 1-3=Moderate Risk; 4-5=High Risk; 6+ =Very High Risk</p>	<p>Before providing care, assess behaviours for potential risk</p> <ul style="list-style-type: none"> <input type="checkbox"/> History of Violence <input type="checkbox"/> Confused <input type="checkbox"/> Irritable <input type="checkbox"/> Boisterous <input type="checkbox"/> Verbal Threats <input type="checkbox"/> Physical Threats <input type="checkbox"/> Agitate/Impulsive <input type="checkbox"/> Paranoid/Suspicious <input type="checkbox"/> Substance <input type="checkbox"/> Intoxication/withdrawal <input type="checkbox"/> Socially appropriate/disruptive <input type="checkbox"/> Body Language <p>Each Check = 1 pt 0=Low Risk; 1-3=Moderate Risk; 4-5=High Risk; 6+ =Very High Risk</p>	<p>Before providing care, assess behaviours for potential risk</p> <ul style="list-style-type: none"> <input type="checkbox"/> History of Violence <input type="checkbox"/> Confused <input type="checkbox"/> Irritable <input type="checkbox"/> Boisterous <input type="checkbox"/> Verbal Threats <input type="checkbox"/> Physical Threats <input type="checkbox"/> Agitate/Impulsive <input type="checkbox"/> Paranoid/Suspicious <input type="checkbox"/> Substance <input type="checkbox"/> Intoxication/withdrawal <input type="checkbox"/> Socially appropriate/disruptive <input type="checkbox"/> Body Language <p>Each Check = 1 pt 0=Low Risk; 1-3=Moderate Risk; 4-5=High Risk; 6+ =Very High Risk</p>	<p>Before providing care, assess behaviours for potential risk</p> <ul style="list-style-type: none"> <input type="checkbox"/> History of Violence <input type="checkbox"/> Confused <input type="checkbox"/> Irritable <input type="checkbox"/> Boisterous <input type="checkbox"/> Verbal Threats <input type="checkbox"/> Physical Threats <input type="checkbox"/> Agitate/Impulsive <input type="checkbox"/> Paranoid/Suspicious <input type="checkbox"/> Substance <input type="checkbox"/> Intoxication/withdrawal <input type="checkbox"/> Socially appropriate/disruptive <input type="checkbox"/> Body Language <p>Each Check = 1 pt 0=Low Risk; 1-3=Moderate Risk; 4-5=High Risk; 6+ =Very High Risk</p>	<p>Before providing care, assess behaviours for potential risk</p> <ul style="list-style-type: none"> <input type="checkbox"/> History of Violence <input type="checkbox"/> Confused <input type="checkbox"/> Irritable <input type="checkbox"/> Boisterous <input type="checkbox"/> Verbal Threats <input type="checkbox"/> Physical Threats <input type="checkbox"/> Agitate/Impulsive <input type="checkbox"/> Paranoid/Suspicious <input type="checkbox"/> Substance <input type="checkbox"/> Intoxication/withdrawal <input type="checkbox"/> Socially appropriate/disruptive <input type="checkbox"/> Body Language <p>Each Check = 1 pt 0=Low Risk; 1-3=Moderate Risk; 4-5=High Risk; 6+ =Very High Risk</p>
--	--	--	--	--



Visit: pshsa.ca/workplace-violence

Poster Instructions

1. Download the poster from www.pshsa.ca/workplace-violence
2. Post in a visible area to remind staff to use the violence assessment tool.

Poster



Visit: pshsa.ca/workplace-violence

Assess the Risk

- ☐ A History of Violence
- ☐ Confused
- ☐ Irritable
- ☐ Boisterous
- ☐ Verbal Threats
- ☐ Physical Threats
- ☐ Agitated/Impulsive
- ☐ Paranoid/Suspicious
- ☐ Substance Intoxication/Withdrawal
- ☐ Socially Inappropriate/Disruptive
- ☐ Defensive Body Language

Rate the Risk

Each Yes = 1 pt. .

- 0 pts =Low Risk;
- 1-3 pts =Moderate Risk;
- 4-5pts =High Risk;
- 6+ pts =Very High Risk

Take Action

- Monitor and remain alert
- Communicate changes in behaviours that may put others at risk
- Initiative violence prevention care planning process
- Apply flag alerts
- Notify manager/supervisor
- Alert Security-assistance may be required
- Use effective therapeutic communication and de-escalation techniques
- Be prepared to apply behavioral management and self-protection techniques
- Initiate appropriate referrals if required
- Ensure communication devices/processes are in place
- Inform Client of VAT results when it is safe to do so
- Call 911 / Initiate Code White Response as necessary

Involve the Client

- Ask the client to help us provide the best possible care by describing known triggers and ways to reduce these behaviours



Your Health. Your Safety. Our Commitment.

© Public Services Health and Safety Association

References

Violence Assessment Tool (VAT)

- Almvik, R., Woods, P., & Rasmussen, K. (2000). Brøset violence checklist: Sensitivity, specificity and interrater reliability. *Journal of Interpersonal Violence*, 15(12), 1284-1296.
- Almvik, R., Woods, P., & Rasmussen, K. (2007). Assessing risk for imminent violence in the elderly: The Brøset violence checklist. *International Journal of Geriatric Psychiatry*, 22, 862-867.
- Canadian Institute for Health Information. (2002). Minimum data set (MDS) 2.0 Canadian version MDS 2.0 form. Ottawa: CIHI.
- Health Care and Residential Facilities Regulation, O. Reg. 67/93. (1990). Retrieved from http://www.e-laws.gov.on.ca/html/regs/english/elaws_regs_930067_e.htm
- Lakeridge Health. (n.d.). Aggressive behaviour scale. Unpublished instrument.
- Manitoba Provincial Healthcare Violence Prevention Program. (n.d.). Screening tool for violence and aggression. Unpublished instrument.
- Marques, M. I., Bessa, A., Santos, L., & Carvalho, S. (2014). Assessment of the risk of violence in Portuguese psychiatric settings using the Brøset violence checklist. In Needham, I., Kingma, M., McKenna, K., Frank, O., Tuttas, C., Kingma, S., & Oud, N. (Ed.), *Fourth International Conference on Violence in the Health Sector* (pp.182). Amsterdam: Kavanah.
- Navarro, J., & Karlins, M. (2008). *What every BODY is saying: An ex-FBI agent's guide to speed-reading people*. New York, NY: HarperCollins Publishers.
- Occupational Health and Safety Act, R.S.O. (1990) c. O.1. Retrieved from http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_90o01_e.htm
- Ogloff, J. R. P., & Daffern, M. (2006). The dynamic appraisal of situational aggression: An instrument to assess for risk for imminent aggression in psychiatric inpatients. *Behavioral Sciences and the Law*, 24, 799-813.
- Public Services Health & Safety Association (n.d.). Completing the violence/aggression assessment checklist (VAAC) for Emergency Departments (ED) or Emergency Medical Services (EMS). Retrieved from <http://www.pshsa.ca/products/violenceaggression-assessment-checklist-vaac/>
- Woods, P., Ashley, C., Kayto, D., & Heusdens, C. (2008). Piloting violence and incident reporting measures on one acute care health inpatient unit. *Issues in Mental Health Nursing*, 29, 455-469.

Sample Interventions

- Advanced Gerontological Education. (2015). Retrieved from <https://www.ageinc.ca/>
- AVP Behavioral Health Services (n.d.). Behavioral health violence risk screening tool. Unpublished instrument.
- Behavioural Supports Ontario. (2012). Behavioural Education and Training Supports Inventory (BETSI). Retrieved from <http://brainxchange.ca/Public/Files/BSO/BETSI-Full-Version-Updated-191212.aspx>
- Crisis Prevention Intervention. (2015). Retrieved from <http://www.crisisprevention.com/>
- CSA Group. (2013). CAN/CSA-Z1003-13/BNQ, 9700-803/2013 - Psychological health and safety in the workplace – prevention, promotion, and guidance to staged implementation. Retrieved from <http://shop.csa.ca/en/canada/occupational-health-and-safety-management/canca-z1003-13bnq-9700-8032013/invnt/z10032013>
- Health Care and Residential Facilities Regulation*, O. Reg. 67/93. (1990). Retrieved from http://www.e-laws.gov.on.ca/html/regis/english/elaws_regs_930067_e.htm
- Mitchell, J. T. (n.d.). Critical incident stress debriefing (CISD). Retrieved from <http://drustvo-psihologa.rs.ba/wp-content/uploads/2014/05/CriticalIncidentStressDebriefing.pdf>
- Navarro, J., & Karlins, M. (2008). *What every BODY is saying: An ex-FBI agent's guide to speed-reading people*. New York, NY: HarperCollins Publishers.
- Occupational Health and Safety Act*, R.S.O. (1990) c. O.1. Retrieved from http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_90o01_e.htm
- Public Services Health & Safety Association. (2006). *A guide to the development of a workplace violence prevention program: Implementing the program in your organization – book 1*. Toronto, ON: Ontario Safety Association for Community & Healthcare.
- Public Services Health & Safety Association. (2006). *A guide to the development of a workplace violence prevention program: Preventing client aggression through gentle persuasive approaches – book 4*. Toronto, ON: Ontario Safety Association for Community & Healthcare.
- Public Services Health & Safety Association. (2009). *Assessing violence in the community: A handbook for the workplace*. Retrieved from <http://www.pshsa.ca/products/assessing-violence-in-the-community-a-handbook-for-the-workplace/>

Individual Client Risk Assessment (ICRA) Toolkit

- Public Services Health & Safety Association. (2011). *Community care: A tool to reduce workplace hazards* (2nd ed.). [Online version]. Retrieved from <https://www.pshsa.ca/products/community-care-a-tool-to-reduce-workplace-hazards-2/>
- Public Services Health & Safety Association. (n.d.). Violence/aggression assessment checklist. Unpublished instrument.
- Safe Management Group. (2014). Retrieved from <http://safemanagement.org/>
- Wiskow, C. (2003). *Guidelines on workplace violence in the health sector. Comparison of major known national guidelines and strategies: United Kingdom, Australia, Sweden, USA (OSHA and California)*. Geneva, Switzerland: ILO/ICN/WHO/PSI. Retrieved from http://www.who.int/violence_injury_prevention/violence/interpersonal/en/WV_ComparisonGuidelines.pdf

Sample Policy

- Almvik, R., Woods, P., & Rasmussen, K. (2000). Brøset violence checklist: Sensitivity, specificity and interrater reliability. *Journal of Interpersonal Violence*, 15(12), 1284-1296.
- Almvik, R., Woods, P., & Rasmussen, K. (2007). Assessing risk for imminent violence in the elderly: The Brøset violence checklist. *International Journal of Geriatric Psychiatry*, 22, 862-867.
- Canadian Institute for Health Information. (2002). Minimum data set (MDS) 2.0 Canadian version MDS 2.0 form. Ottawa: Canadian Institute for Health Information.
- Gormley, N., Lyons, D., & Howard, R. (2001). Behavioural management of aggression in dementia: A randomized controlled trial. *Age and Ageing*, 30(2), 141-145.
- Health Care and Residential Facilities Regulation*, O. Reg. 67/93. Retrieved from http://www.e-laws.gov.on.ca/html/regs/english/elaws_regs_930067_e.htm
- Lakeridge Health. (n.d.). Aggressive behaviour scale. Unpublished instrument.
- Lundrigan, E., Hutchings, D., Mathews, M., Lynch, A., & Goosney, J. (2010). A risk assessment screening tool for community health care workers. *Home Health Care Management & Practice*, 22(6), 403-407.
- Manitoba Provincial Healthcare Violence Prevention Program. (n.d.). Screening tool for violence and aggression. Unpublished instrument.
- Marques, M. I., Bessa, A., Santos, L., & Carvalho, S. (2014). Assessment of the risk of violence in Portuguese psychiatric settings using the Brøset violence checklist. In Needham, I., Kingma, M., McKenna, K., Frank, O., Tuttas, C., Kingma, S., & Oud, N. (Ed.), *Fourth International Conference on Violence in the Health Sector* (pp.182). Amsterdam: Kavanah.

Individual Client Risk Assessment (ICRA) Toolkit

Navarro, J., & Karlins, M. (2008). *What every BODY is saying: An ex-FBI agent's guide to speed-reading people*. New York, NY: HarperCollins Publishers.

Ogloff, J. R. P., & Daffern, M. (2006). The dynamic appraisal of situational aggression: An instrument to assess for risk for imminent aggression in psychiatric inpatients. *Behavioral Sciences and the Law*, 24, 799-813.

Occupational Health and Safety Act, R.S.O. (1990) c. O.1. Retrieved from http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_90o01_e.htm

Public Services Health & Safety Association. (2006). *A guide to the development of a workplace violence prevention program: Implementing the program in your organization – book 1*. Toronto, ON: Ontario Safety Association for Community & Healthcare.

Public Services Health & Safety Association. (2006). *A guide to the development of a workplace violence prevention program: Preventing client aggression through gentle persuasive approaches – book 4*. Toronto, ON: Ontario Safety Association for Community & Healthcare.

Public Services Health & Safety Association. (2009). *Assessing violence in the community: A handbook for the workplace*. Retrieved from <http://www.pshsa.ca/products/assessing-violence-in-the-community-a-handbook-for-the-workplace/>

Public Services Health & Safety Association. (2011). *Community care: A tool to reduce workplace hazards* (2nd ed.). [Online version]. Retrieved from <https://www.pshsa.ca/products/community-care-a-tool-to-reduce-workplace-hazards-2/>

Wiskow, C. (2003). *Guidelines on workplace violence in the health sector. Comparison of major known national guidelines and strategies: United Kingdom, Australia, Sweden, USA (OSHA and California)*. Geneva, Switzerland: ILO/ICN/WHO/PSI. Retrieved from http://www.who.int/violence_injury_prevention/violence/interpersonal/en/WV_ComparisonGuidelines.pdf



Individual Client Risk Assessment Toolkit for Health Care Settings

Public Services Health and Safety Association (PSHSA)

4950 Yonge Street, Suite 1800

Toronto, Ontario M2N 6K1

Canada

Telephone: 416-250-2131

Fax: 416-250-7484

Toll Free: 1-877-250-7444

Web site: www.pshsa.ca

Connect with us:

 [@PSHSAca](https://twitter.com/PSHSAca)

Product Code: VPRASEEN0417